

Graded q -pseudo-differential Operators and Supersymmetric Algebras

Ahmed Jellal^{a *} and El Hassan El Kinani^{b †}

^a *High Energy Physics Section*

the Abdus Salam International Centre for Theoretical Physics

Strada Costiera 11, 34100 Trieste, Italy

^b *Mathematical Physics Group, Dept. of Mathematics*

Faculty of Sciences and Technics, University Molay Ismail

P.O.B. 509, Errachidia, Morocco

Abstract

We give a supersymmetric generalization of the sine algebra and the quantum algebra $U_t(sl(2))$. Making use of the q -pseudo-differential operators graded with a fermionic algebra, we obtain a supersymmetric extension of sine algebra. With this scheme we also get a quantum superalgebra $U_t(sl(2/1))$.

PACS: 02.20, 03.65

Keywords: Graded q -pseudo-differential operators, supersymmetric sine algebra, quantum superalgebra

*E-mail: jellal@ictp.trieste.it – jellal@youpy.co.uk

†E-mail: el_kinani@fste.ac.ma

1 Introduction

One of the most important infinite-dimensional Lie algebra is the one generated by the so-called pseudo-differential operators [1]. This can be viewed as a generalization of the Virasoro algebra and of the Lie algebra of differential operators. Recently, the supersymmetric algebras have been applied to the study of some physical problems. For example, the supersymmetric sine algebra is used to investigate the properties of Bloch electron in a constant uniform magnetic field [2]. Moreover, the quantum superalgebras are applied to solve some problems for instance related to superconductivity [3] and the quantum Hall effect [4]. These results lead us to think to the present work.

In this paper, we will present an approach to obtain a realization of certain supersymmetric algebras. More precisely, we will propose a graded q -pseudo-differential operator realization of the supersymmetric extension of the sine algebra and the quantum superalgebra $U_t(sl(2/1))$.

This paper is organized as follows: In section 2 we review some basic notions related to q -pseudo-differential operators and also to the realization of sine algebra and $U_t(sl(2))$ in this framework. We propose a realization of the supersymmetric extension of the last algebras in section 3. We give a conclusion in the final section.

2 Preliminaries

Before going on, we would like to give a short review concerning some basic notions, which will be useful in the next section. This concerns the q -pseudo-differential operators, sine algebra and $U_t(sl(2))$.

2.1 q -pseudo-differential operators

We start first by defining the so-called q -derivation. For this matter, let q be a complex number different from 0 and 1. By definition, the q -derivation or more generally α -

derivation is given by

$$d_\alpha(fg) = \alpha(f)d_\alpha(g) + d_\alpha(f)g, \quad (1)$$

where $f, g \in C[x, x^{-1}]$ are the ring of polynomials in an indeterminant x and its inverse x^{-1} . In eq.(1), α is a linear mapping. An example of α -derivation is given by Jackson's q -differential operator ∂_q , such as [5]

$$\partial_q(f) = \frac{f(qx) - f(x)}{(q-1)x}, \quad (2)$$

which leads to the following form for eq.(1)

$$\partial_q(fg) = \eta_q(f)\partial_q(g) + \partial_q(f)g, \quad (3)$$

where the shift operator η_q is

$$\eta_q(f(x)) = f(qx). \quad (4)$$

Now let us introduce the q -pseudo-differential operators algebra $q - \psi DO$. The latter is characterized by the relation [5]

$$q - \psi DO = P(x, \partial_q) = \sum_{i=-\infty}^N P_i(x) \partial_q^i, \quad P_i(x) \in C[x, x^{-1}]. \quad (5)$$

Consequently, the algebra $q - \psi DO$ is generated by $x, x^{-1}, \partial_q, \partial_q^{-1}$ with the relation

$$\partial_q x - qx \partial_q = 1. \quad (6)$$

Note that the family $\{x^i \partial_q^j\}_{i,j \in \mathbb{Z}}$ forms a basis of $q - \psi DO$. Then, the algebra $q - \psi DO$ can be viewed as a Lie algebra, which can be defined by the commutation relation

$$[P, Q] = P \circ Q - Q \circ P, \quad (7)$$

where the multiplication law "o" is

$$\begin{aligned} \partial_q \circ f &= \eta_q(f) \partial_q + \partial_q f, \\ \partial_q^{-1} \circ f &= \sum_{k \geq 0} (-1)^k q^{-k(k+1)/2} (\eta_q^{-k-1}(\partial_q^k f)) \partial_q^{-k-1}. \end{aligned} \quad (8)$$

The last equation are obtained by using the following relation

$$\partial_q^{-1} \circ \partial_q \circ f = \partial_q \circ \partial_q^{-1} \circ f = f. \quad (9)$$

Note that eq.(8) can be unified as follows

$$\partial_q^n \circ f = \sum_{k \geq 0} \binom{n}{k}_q (\eta_q^{n-k} (\partial_q^k f) \partial_q^{n-k}), \quad (10)$$

for all n . In the last equation, the q -binomials take the form

$$\binom{n}{k}_q = \frac{(n)_q (n-1)_q \dots (n-k+1)_q}{(1)_q (1)_q \dots (k)_q}, \quad (11)$$

and the q -numbers are given by

$$(n)_q = \frac{q^n - 1}{q - 1}, \quad (12)$$

where the convention

$$\binom{n}{0}_q = 1, \quad (13)$$

is taken. We also add that the residue of the symbol $P(x, \partial_q)$ can be written as

$$\mathbf{Res}(\sum_{i=-\infty}^N P_i(x) \partial_q^i) = P_{-1}(x), \quad (14)$$

and its **Tr**-functional is

$$\mathbf{Tr}(\sum_{i=-\infty}^N P_i(x) \partial_q^i) = \int_{s^1} P_{-1}(x) dx. \quad (15)$$

Considering a subfamily of $q - \psi DO$ as

$$q - S\psi DO = \{P(x, \partial_q) = \sum_{i=-\infty}^N P_i(x) \cdot (\eta_q)^i / \text{Tr}(P) = 0\}. \quad (16)$$

From eq.(6), we obtain the relation

$$\eta_q x = qx \eta_q. \quad (17)$$

Therefore, η_q and x generate a non-commutative algebra, which is homomorphic with the Manin's plane "quantum plane" [6].

2.2 Sine algebra

In this subsection, we review the realization of the sine algebra and the quantum algebra $U_t(sl(2))$. To do this, let us consider a subfamily of $q - S\psi DO$ generated by $J_{\mathbf{m}}$. This can be constructed as follows [7]

$$J_{\mathbf{m}} = q^{-m_1 \cdot m_2 / 2} \eta_q^{m_1} x^{m_2}, \quad (18)$$

with $\mathbf{m} = (m_1, m_2)$. Calculating the commutation relation, it is found

$$[J_{\mathbf{m}}, J_{\mathbf{n}}] = (q^{(\mathbf{m} \times \mathbf{n})/2} - q^{-(\mathbf{m} \times \mathbf{n})/2}) J_{\mathbf{m} + \mathbf{n}}, \quad (19)$$

where $\mathbf{m} \times \mathbf{n} = m_1 n_2 - m_2 n_1$. It is interesting to note that when q is a F -th root of unity, i.e. $q = \exp(\frac{4\pi i}{F})$, the last equation takes the following form

$$[J_{\mathbf{m}}, J_{\mathbf{n}}] = 2i \sin(\frac{2\pi}{F} \mathbf{m} \times \mathbf{n}) J_{\mathbf{m} + \mathbf{n}}, \quad (20)$$

which generates the sine algebra or Fairlie-Fletcher-Zachos (FFZ) [8] algebra. This is exactly the Moyal bracket quantization of the area-preserving diffeomorphisms or symplectomorphisms algebra on $2 - d$ torus. It should be mentioned that the deformation here (eq.(20)) is the Moyal quantization which is strongly different from the Drinfel'd and Jimbo one [9], where the Hopf structure plays a crucial role.

Now let us give a construction of the quantum algebra $U_t(sl(2))$ in this scheme. Before going on, we recall that this algebra is defined by the commutation relations [9]

$$\begin{aligned} [X^+, X^-] &= \frac{t^{2h} - t^{-2h}}{t - t^{-1}}, \\ [h, X^{\pm}] &= \pm X^{\pm}, \end{aligned} \quad (21)$$

where t is the deformed parameter, $t \neq 0, 1$. In the limit where $t \rightarrow 1$, the above equations reduce to ones defining the Lie algebra $sl(2)$.

The generators of $U_t(sl(2))$ can be embedded as follows [7]

$$\begin{aligned} X^+ &= \frac{J_{\mathbf{m}} - J_{\mathbf{n}}}{t - t^{-1}}, & X^- &= \frac{J_{-\mathbf{m}} - J_{-\mathbf{n}}}{t - t^{-1}}, \\ t^{+2h} &= J_{\mathbf{m} - \mathbf{n}}, & t^{-2h} &= J_{\mathbf{n} - \mathbf{m}}. \end{aligned} \quad (22)$$

they satisfy the commutation relations given by eq.(21) if t is given by

$$t = q^{\mathbf{m} \times \mathbf{n}/2}. \quad (23)$$

This concludes a short review of the q -pseudo-differential operators and its related algebras. Now let us address our main goal, which will be expanded in the next sections.

3 Supersymmetric extension

In this section we start our generalization of the above results. Otherwise, we are looking for a supersymmetric extension of the sine algebra and the quantum algebra $U_t(sl(2))$. To do this let us begin by the realization of the supersymmetric sine algebra. This task is the subject of the following subsection.

3.1 Supersymmetric sine algebra

Our goal here is to extend the sine algebra to supersymmetric case. For this matter, let us introduce a fermionic algebra. We begin by considering the following matrices [10]

$$f = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad f^+ = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad (24)$$

it is easy to show that the generators f and f^+ expanding a fermionic algebra. The latter is characterized by the relations

$$ff^+ + f^+f = 1, \quad f^2 = 0 = (f^+)^2. \quad (25)$$

To obtain the supersymmetric extension of the operators $J_{\mathbf{m}}$, we can proceed as follows

$$\begin{aligned} T_{\mathbf{m}} &= J_{\mathbf{m}} \otimes (ff^+ + f^+f), \\ S_{\mathbf{m}} &= J_{\mathbf{m}} \otimes (ff^+ - f^+f), \end{aligned} \quad (26)$$

which define the so-called graded q -pseudo-differential operators. These operators can be written as

$$T_{\mathbf{m}} = J_{\mathbf{m}} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad S_{\mathbf{m}} = J_{\mathbf{m}} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (27)$$

Using the last equation, we prove that the operators $T_{\mathbf{m}}$ and $S_{\mathbf{m}}$ satisfy the following relations

$$\begin{aligned} [T_{\mathbf{m}}, T_{\mathbf{n}}]_- &= (q^{(\mathbf{m} \times \mathbf{n})/2} - q^{-(\mathbf{m} \times \mathbf{n})/2}) T_{\mathbf{m}+\mathbf{n}}, \\ [T_{\mathbf{m}}, S_{\mathbf{n}}]_- &= (q^{(\mathbf{m} \times \mathbf{n})/2} - q^{-(\mathbf{m} \times \mathbf{n})/2}) S_{\mathbf{m}+\mathbf{n}}, \\ [S_{\mathbf{m}}, S_{\mathbf{n}}]_+ &= (q^{(\mathbf{m} \times \mathbf{n})/2} + q^{-(\mathbf{m} \times \mathbf{n})/2}) T_{\mathbf{m}+\mathbf{n}}. \end{aligned} \quad (28)$$

Let us remember that our goal here is to obtain a generalization of the sine algebra given by eq.(20). For this matter, we consider a q F th root of unity. In this case, we show that the relations are verified

$$\begin{aligned} [T_{\mathbf{m}}, T_{\mathbf{n}}]_- &= 2i \sin(\frac{2\pi}{F}(\mathbf{m} \times \mathbf{n})) T_{\mathbf{m}+\mathbf{n}}, \\ [T_{\mathbf{m}}, S_{\mathbf{n}}]_- &= 2i \sin(\frac{2\pi}{F}(\mathbf{m} \times \mathbf{n})) S_{\mathbf{m}+\mathbf{n}}, \\ [S_{\mathbf{m}}, S_{\mathbf{n}}]_+ &= 2 \cos(\frac{2\pi}{F}(\mathbf{m} \times \mathbf{n})) T_{\mathbf{m}+\mathbf{n}}. \end{aligned} \quad (29)$$

This set of equations generate exactly the supersymmetric sine algebra [11]. In the next section we show how to obtain $U_t(sl(2/1))$.

3.2 Quantum superalgebra $U_t(sl(2/1))$

The task of the present section is to realize the quantum superalgebra $U_t(sl(2/1))$ based on the defined supersymmetry operators Eq.(26). We begin by recalling that the quantum superalgebra $U_t(sl(2/1))$ can be viewed as a t -deformation of the classical Lie superalgebra $sl(2/1)$ through the t -deformed relations, between a set of generators denoted by $e_1, e_2, f_1, f_2, k_1 = t^{h_1}, k_1^{-1} = t^{-h_1}, k_2 = t^{h_2}$ and $k_2^{-1} = t^{-h_2}$, such as [12]

$$\begin{aligned} k_1 k_2 &= k_2 k_1, & k_i e_j k_i^{-1} &= t^{a_{ij}} e_j, & k_i f_j k_i^{-1} &= t^{-a_{ij}} f_j, \\ e_1 f_1 - f_1 e_1 &= \frac{k_1^2 - k_1^{-2}}{t - t^{-1}}, & e_2 f_2 + f_2 e_2 &= \frac{k_2^2 - k_2^{-2}}{t - t^{-1}}, \\ e_1 f_2 - f_2 e_1 &= 0, & e_2 f_1 - f_1 e_2 &= 0, & e_2^2 &= 0 = f_2^2, \\ e_1^2 e_2 - (t + t^{-1}) e_1 e_2 e_1 + e_2 e_1^2 &= 0, & f_1^2 f_2 - (t + t^{-1}) f_1 f_2 f_1 + f_2 f_1^2 &= 0. \end{aligned} \quad (30)$$

The last two relations are called the Serre relations. The matrix (a_{ij}) is the Cartan one of $sl(2/1)$, i.e.

$$(a_{ij}) = \begin{pmatrix} 2 & -1 \\ -1 & 0 \end{pmatrix}. \quad (31)$$

$U_t(sl(2/1))$ is a quasi-triangular Hopf superalgebra endowed with the \mathbf{Z}_2 -graded Hopf algebra structure

$$\begin{aligned}\Delta(k_i) &= k_i \otimes k_i, & \Delta(e_i) &= e_i \otimes \mathbf{1} + e_i \otimes k_i, & \Delta(f_i) &= e_i \otimes k_i^{-1} + \mathbf{1} \otimes f_i, \\ \epsilon(k_i) &= \mathbf{1}, & \epsilon(e_i) &= \epsilon(f_i) = 0, \\ S(e_i) &= -k_i^{-1}e_i, & S(e_i) &= -f_i k_i & S(k_i^{\pm 1}) &= k_i^{\pm 1}, \quad i = 1, 2.\end{aligned}\tag{32}$$

The \mathbf{Z}_2 -grading of the generators are $[e_2] = [f_2] = 1$ and zero otherwise. The multiplication rule for the tensor product is \mathbf{Z}_2 -graded and is defined for the elements a, b, c, d of $U_t(sl(2/1))$ by

$$(a \otimes b)(c \otimes d) = (-1)^{[b][c]}(ac \otimes bd).\tag{33}$$

With the help of the symmetry operators Eq.(26), it is possible to give the following construction for the generators e_1, e_2, f_1, f_2, k_1 and k_2

$$\begin{aligned}e_1 &= \frac{T_{(m_1, m_2)} + T_{(-m_1, m_2)}}{t - t^{-1}} \otimes \mathbf{1}, & f_1 &= -i \frac{T_{(m_1, -m_2)} - T_{(-m_1, -m_2)}}{t - t^{-1}} \otimes \mathbf{1}, \\ k_1 &= -iT_{(-2m_1, 0)} \otimes \mathbf{1}, & k_1^{-1} &= iT_{(2m_1, 0)} \otimes \mathbf{1}, \\ k_2 &= -iT_{(-2m_1, 0)} \otimes \begin{pmatrix} t^{-2} & 0 \\ 0 & t^2 \end{pmatrix}, & k_2^{-1} &= iT_{(2m_1, 0)} \otimes \begin{pmatrix} t^2 & 0 \\ 0 & t^{-2} \end{pmatrix}, \\ e_2 &= \frac{T_{(m_1, -m_2)} - T_{(-m_1, -m_2)}}{(t - t^{-1})^{\frac{1}{2}}} \otimes f, & f_2 &= \frac{T_{(m_1, m_2)} + T_{(-m_1, m_2)}}{(t - t^{-1})^{\frac{1}{2}}} \otimes f^+.\end{aligned}\tag{34}$$

It turns out that the above generators satisfy the algebraic relations characterizing the quantum superalgebra $U_t(sl(2/1))$ as shown by Eq.(30) where the t -deformed parameter is given by

$$t = e^{m_1 m_2}.\tag{35}$$

This is a way to prove that we can realize the $U_t(sl(2/1))$ by using the supersymmetry operators stated above.

4 Conclusion

In this paper, we have shown how the graded q -pseudo-differential operators lead to obtain a supersymmetric extension of the sine algebra and the quantum algebra $U_t(sl(2))$. Otherwise, we have realized the the supersymmetric sine algebra and the quantum superalgebra $U_t(sl(2/1))$ with the help of q -operators graded with a fermionic algebra.

Acknowledgement

A. Jellal wishes to thank Prof. S. Randjbar-Daemi, Head of High Energy Section of the Abdus Salam International Centre for Theoretical Physics (AS-ICTP), for the kind hospitality at his section. He would like also to express his gratitude to Prof. E.H. Saidi for his encouragement. The authors are grateful to Prof. G. Thompson for reading the manuscript and A. Jellal acknowledges him for his kind encouragement. A. Jellal is thankful to Prof. C.K. Zachos for drawing his attention to reference [11].

References

- [1] I.M. Gel'fand and L.A. Dickey, *Funct. Ann. App.* **10** (1976)252; V.G. Kac et al, **MIT/27/87**; A.O. Radul, *Phys. Lett.* **B265** (1991)86; C. Kassel, *Commun. Math. Phys.* **140** (1992)343; B. Khesin, V. Lyubashenko and C. Roger, **hep-th/9403189**.
- [2] T. Dereli and A. Verçin, *J. Phys. A: Math. Gen* **26** (1993)6961; *ibid*, *Phys. Lett.* **B288** (1992)109.
- [3] M.D. Gould, K.E. Hibberd and J.R. Links, *Phys. Lett.* **A212** (1996)156.
- [4] A. Jellal, *Rep. Math. Phys.* **47** (2000), in press.
- [5] C. Kassel, *Commun. Math. Phys.* **140** (1992)343.
- [6] Yu. Manin, *Commun. Math. Phys.* **123** (1989)163.
- [7] E.H El Kinani, *Mod. Phys. Lett.* **A14** (1999)1609.
- [8] D.B. Fairlie, P. Flecher and C. Zachos, *J. Math. Phys.* **31** (1990)1088.
- [9] V.G. Drinfeld, *Proc. Int. Congr. Math.* ed A.M. Gleason (Berkeley: MSRI 1986) p798; M. Jimbo, *Lett. Math. Phys.* **10** (1985)63; *ibid Lett. Math. Phys.* **11** (1986)247.
- [10] A. Jellal, *Mod. Phys. Lett.* **A14** (1999)2253.

- [11] D.B. Fairlie and C.K. Zachos, *Phys. Lett.* **B224** (1989)101.
- [12] M. Chaichian and P.P. Kulish, *Phys. Lett.* **B234** (1990)72; B. Abdesselam, D. Arnaudon and M. Bauer, *J. Phys. A: Math. Gen* **30** (1997)867.